

# End-of-Life Care: Beyond Survival, Towards Dignity

– Dr. Yash Javeri<sup>1</sup> 

<sup>1</sup> Director, Critical Care and Emergency Medicine, Regency Super Specialty Hospital, Lucknow, Uttar Pradesh, India.  
E mail for correspondence- dryashjaveri@yahoo.com

## Introduction

Modern medicine has transformed the boundaries of human survival. Advances in intensive care, mechanical ventilation, renal replacement therapy, vasopressor support, artificial nutrition, cardiac devices, antimicrobials, and organ support technologies have allowed clinicians to sustain life in situations that were once considered irreversible. These developments are among the greatest achievements of medical science. They have saved countless lives, restored health, and extended meaningful survival for many patients. Yet, these same advances have also created new ethical and clinical dilemmas. Medicine can now maintain biological function even when the possibility of meaningful recovery has disappeared. A heart may continue to beat, lungs may be ventilated, blood pressure may be supported, and nutrition may be delivered artificially, but the patient may no longer have awareness, interaction, comfort, or any realistic chance of recovery.

This reality compels medicine to ask a difficult but necessary question: when does sustaining life cease to be healing and begin to prolong dying? This question is not merely theoretical. It is faced daily in emergency departments, intensive care units, oncology wards, neurology units, and homes where families struggle to decide what is right for their loved ones. End-of-life care begins at this intersection. It asks clinicians, families, institutions, and society to recognize that the purpose of medicine is not merely to extend life at all costs. The purpose of medicine is also to relieve suffering, respect human values, and preserve dignity when cure is no longer possible [1-3].

## The Meaning of Dignity at the End of Life

Dignity is often discussed as a philosophical or moral ideal, but in end-of-life care it is also a clinical responsibility. A dignified death does not mean abandoning treatment. It means ensuring that the patient is not reduced to a collection of failing organs, laboratory values, ventilator settings, and drug infusions. Dignity requires that the patient remains at the centre of care. Their comfort, values, beliefs, previously expressed wishes, and emotional needs must guide decision-making. The clinical team must ask not only “Can we continue treatment?” but also “Should we continue this treatment?” and “Is this treatment still serving the patient?” Preserving dignity also means protecting the patient from unnecessary pain, invasive procedures, isolation, and loss of privacy [2,4,5]. It requires the treating team to recognize

the patient as a person with a life story, relationships, fears, hopes, and values. Even when consciousness is lost, the patient's dignity must remain fully respected.

When recovery becomes impossible, the goal of care must shift deliberately and responsibly. The focus moves from prolonging life to preserving dignity, from intervention to intention, from survival alone to comfort and meaning, from aggressive treatment to compassionate care, and from doing everything possible to doing everything appropriate. This transition is not a failure of medicine. It is a sign of mature, ethical, and patient-centred medical practice. It reflects the understanding that medical success cannot be measured only by survival statistics, but also by the quality of care delivered in the final phase of life. Such a shift requires courage from clinicians and trust from families. It may be emotionally difficult because it involves accepting the limits of treatment. However, when done with honesty and compassion, it allows the patient's remaining time to be protected from avoidable suffering.

## Life-Sustaining Treatments are not a Moral Obligation

Life-sustaining treatments include mechanical ventilation, vasopressor support, dialysis, cardiopulmonary resuscitation, invasive monitoring, antibiotics, blood products, clinically assisted nutrition, and hydration. These interventions can be lifesaving when used in reversible illness. However, they are still medical treatments. Like all treatments, they must be evaluated in relation to benefit, burden, prognosis, and alignment with the patient's goals.

No treatment is automatically obligatory simply because it is available. The mere ability to continue an intervention does not make its continuation ethically necessary [6,7]. Modern medicine must therefore distinguish between treatments that restore meaningful life and treatments that merely delay death. This distinction is especially important in intensive care settings, where machines and drugs may create the appearance of stability while the underlying disease remains irreversible. Families may interpret continued treatment as continued hope. It is the responsibility of clinicians to explain when treatment is no longer changing the outcome.

A treatment is appropriate when its expected benefit outweighs its burden. However, when treatment no longer offers meaningful recovery, symptom relief, or improvement in quality of life, it may become disproportionate [8-10]. Continuing such treatment can increase suffering rather than reduce it. For example,

mechanical ventilation may be appropriate in a patient with reversible respiratory failure. But in a patient with irreversible neurological injury, progressive multi-organ failure, or terminal illness with no possibility of recovery, ventilation may only prolong the dying process. Similarly, vasopressors, dialysis, and artificial feeding may be beneficial in selected situations, but they may become medically inappropriate when they only sustain organ function without improving the patient's overall condition or comfort. Proportionality requires a balanced view of what the treatment is achieving and what it is costing the patient physically, emotionally, and spiritually.

The burden of treatment may include pain, repeated procedures, loss of communication, prolonged ICU stay, distress to families, and loss of a peaceful dying process [11]. Therefore, benefit must not be assessed in purely physiological terms. It must be assessed in terms of the patient's overall welfare. Clinically assisted nutrition and hydration are often emotionally perceived as basic care [12-15]. Families may feel that stopping artificial feeding is equivalent to starving the patient. However, in medical and ethical terms, artificial nutrition and hydration are medical interventions. They require tubes, devices, monitoring, and clinical judgment.

When they no longer serve the patient's interests, or when they contribute to discomfort, complications, aspiration, fluid overload, or prolongation of dying, they may be ethically withheld or withdrawn. This is particularly relevant in irreversible conditions such as persistent vegetative state, advanced dementia, terminal illness, or severe irreversible brain injury. It is important to explain this distinction gently to families. Food and water have deep emotional and cultural meaning, and decisions about artificial nutrition can be painful. Clinicians must reassure families that comfort measures, mouth care, symptom relief, and compassionate presence will continue even if medically assisted feeding is not pursued.

### **Futility in End-of-Life Care**

Futility refers to treatment that is unlikely to achieve its intended physiological or meaningful clinical goal [16-19]. It does not mean that the patient's life is futile. It means that a particular medical intervention is no longer useful for achieving recovery, comfort, or meaningful survival. This distinction is extremely important. The dignity of the patient must never be questioned. What may be questioned is the usefulness of continued intervention. The term futility should therefore be used carefully and respectfully. Families may misunderstand it as a judgment on the value of their loved one's life. Clinicians should instead explain that the disease has reached a stage where further intervention cannot reverse the condition or restore meaningful recovery.

The decision that treatment is futile must be based on careful medical assessment. It should not arise from emotional exhaustion, moral distress, bed shortages, financial constraints, or institutional pressure [9,10, 19-23]. Clinical futility must be judged through reversibility of the underlying illness, prognosis, response to ongoing treatment, expected quality of life, patient's known values and preferences, consensus among treating clinicians, and multidisciplinary discussion when needed. Such decisions should be transparent, documented, and communicated compassionately to the family. The process should allow time for questions, second opinions when appropriate, and repeated discussions if the family is struggling to understand the situation. Objective judgment also protects clinicians from making decisions based on impulse or fatigue. End-of-life decisions are too serious to be made casually. They require clinical clarity, ethical reflection, and institutional support.

The concept of futility must be used with great caution. It should not become a convenient label for difficult cases. It should not be used to deny care to patients who are elderly, disabled, economically disadvantaged, or socially vulnerable. Futility must remain anchored in medical reasoning, ethical reflection, and respect for the patient. The patient's age, disability, social status, or economic background must never become the basis for limiting care. Every patient deserves individualized assessment. A treatment that is inappropriate for one patient may be appropriate for another depending on prognosis, reversibility, and the patient's own goals. Ethical decision-making must therefore remain patient-specific and free from prejudice.

### **Withholding and Withdrawing Treatment**

Withholding treatment means deciding not to start a medical intervention because it is unlikely to benefit the patient [9, 10, 24, 25]. Withdrawing treatment means stopping an intervention that has already been started because it is no longer beneficial [24-27]. Emotionally, withdrawing treatment often feels more difficult than withholding it. Families and clinicians may feel that stopping a ventilator, dialysis, or vasopressor support is an active step. However, ethically, withholding and withdrawing inappropriate treatment are similar when the treatment no longer serves the patient's goals. This understanding is essential because many treatments are started during uncertainty or emergency. Once more information becomes available and prognosis becomes clearer, it is ethically acceptable to reassess and discontinue treatments that are no longer useful. Starting treatment should not mean that it must continue indefinitely.

A critical ethical distinction must be preserved: withholding or withdrawing life-sustaining treatment is not the same as

causing death. In such situations, the underlying illness causes death. The physician does not kill the patient. The physician recognizes the limits of medicine and allows the natural course of disease to proceed while ensuring that the patient is comfortable and cared for.

“Care does not stop when treatment stops.

We may withdraw machines, but we never withdraw care.

When cure is no longer possible, comfort becomes the priority.”

This message must be clearly communicated to families. Many relatives fear that limiting treatment means the patient will be neglected [28]. The healthcare team must reassure them that attention, monitoring, symptom relief, nursing care, and emotional support will continue with even greater focus.

When life-sustaining treatment is withheld or withdrawn, the responsibility of the healthcare team increases rather than decreases. The patient must receive meticulous care focused on comfort, relief of suffering, privacy, communication, and emotional support. This includes control of pain and breathlessness, relief of agitation, management of secretions, skin care, nursing comfort, family presence, spiritual support, and clear explanation of what to expect during the dying process.

This is not passive care. It is active, compassionate, and deeply humane care. The aim is to ensure that the patient does not experience avoidable distress and that the family does not feel abandoned. Good end-of-life care also prepares the family for visible changes during dying, such as altered breathing, reduced consciousness, cold extremities, or decreased urine output. Explaining these changes reduces fear and prevents unnecessary panic-driven interventions.

**The Indian Legal and Ethical Context**

In India, end-of-life care has gradually gained ethical and legal

recognition. Landmark judicial decisions have acknowledged that the right to life includes the right to live and die with dignity. These developments have helped clarify the permissibility of withholding and withdrawing life-sustaining treatment under appropriate safeguards. This legal evolution is important because many clinicians in India have historically feared legal consequences when limiting futile treatment (Figure 1). As a result, aggressive interventions have often continued even when they no longer benefit the patient.



**FIGURE-1: End of life care journey of India**

Abbreviations: EOL: End-of-life; ISCCM: Indian Society of Critical Care Medicine; IAPC: Indian Association of Palliative Care; ICMR: Indian Council of Medical Research; DNAR: Do Not Attempt Resuscitation; ICU: Intensive Care Unit; ICUs: Intensive Care Units.

The legal progress also reflects a broader moral recognition that dignity at the end of life is not a luxury. It is a fundamental component of humane care. However, awareness of these developments remains uneven across institutions and among healthcare workers.

Legal frameworks should not intimidate doctors. They should enable ethical care. Clear legal protection is essential for clinicians who act in good faith, follow due process, involve families, document decisions, and prioritize the patient's best interests. However, legal recognition alone is not enough. Hospitals must develop practical systems so that these principles can be implemented at the bedside. The law should provide confidence, not confusion. If procedures are too complex, clinicians may avoid appropriate end-of-life decisions despite legal permission. Therefore, legal clarity must be translated into simple institutional pathways. The aim should be to make ethical end-of-life care a routine part of good clinical practice, not an exceptional act [29].

### **Role of all Stakeholders**

#### *Role of Physicians*

Doctors must not remain passive observers in end-of-life decision-making. They must lead with clinical clarity, ethical sensitivity, and compassionate communication. Physicians should initiate discussions when treatment is no longer beneficial. They should explain prognosis honestly and guide families through difficult decisions. Avoiding these conversations does not protect families. It burdens them with uncertainty and guilt.

The physician's duty does not end when cure is no longer possible. In many ways, the physician's role becomes even more important. Patients and families need guidance, reassurance, and presence. Being present at the end of life is a powerful form of care. Even when no curative treatment remains, the physician can still relieve symptoms, reduce fear, support the family, and ensure that the patient's final phase is peaceful and dignified.

#### *Role of Hospitals and Institutions*

Every hospital should have a clear end-of-life care policy. Such policies should define processes for identifying medically inappropriate treatment, conducting family meetings, documenting decisions, obtaining second opinions, involving ethics committees, and ensuring comfort care.

Without institutional support, clinicians may feel isolated and vulnerable. Institutional policies also create consistency. They prevent end-of-life decisions from varying widely depending on the individual doctor on duty. A clear policy reassures patients, families, and healthcare workers that decisions are being made through a fair and transparent process.

#### *Role of Nurses and Allied Healthcare Workers*

Nurses are often closest to patients and families. They observe suffering, provide comfort, explain bedside changes, and support families during the dying process. Their role in end-of-life care is central. They should be included in decision-making conversations and trained in symptom recognition, comfort care, communication, and family support. Families often trust nurses because of their continuous presence at the bedside. Nurses can recognize subtle signs of discomfort and alert physicians early. Their involvement makes end-of-life care more responsive, compassionate, and practical.

#### *Role of Primary Care Doctors*

Primary care doctors and family physicians are often the first point of contact. They know the patient's family, social background, and long-term illness trajectory. They are well placed to initiate advance care planning and guide families before crisis occurs. Training in basic palliative care and end-of-life communication should be integrated into undergraduate and postgraduate medical education [30-32].

Primary care doctors can also help identify patients who need early palliative support. They can prevent unnecessary hospital visits by managing symptoms locally when appropriate. Their role is especially important for elderly patients, chronically ill patients, and those with limited access to tertiary care.

#### *Role of Community Health Workers*

Community health workers, including ASHA workers, can play an important role in bridging hospitals and homes. They often understand local culture, family dynamics, and community beliefs. With appropriate training, they can help identify patients needing palliative support, guide families, assist with symptom reporting, and encourage timely medical consultation.

They can also help families understand basic comfort measures and when to seek medical help. Their presence can reduce isolation for families caring for dying patients at home. In resource-limited settings, they may become the most accessible link between the healthcare system and the patient.

#### *Role of Society and Culture*

Death remains a taboo subject in many families. It is often discussed only when crisis has already occurred. This silence prevents planning and increases suffering. Society must become more comfortable discussing serious illness, dying, and medical limits [33]. Such conversations do not invite death. They prepare families to face it with clarity and dignity. Schools, communities, religious groups, media, and healthcare institutions can all help normalize these conversations. A society that can speak openly about dying is better able to care for the dying. Silence often leads to

fear; conversation can create preparedness.

#### *Role of Lawmakers and Policymakers*

Lawmakers have a crucial role in bridging the gap between ethical principle and clinical practice. Legal procedures for withholding and withdrawing life-sustaining treatment must be clear, accessible, and practical. Complicated processes may discourage implementation and leave patients exposed to prolonged suffering.

Legal pathways must be understandable for clinicians working in busy hospitals. They should not require excessive paperwork, repeated approvals, or unclear authority. Simplicity is not the enemy of safety; it is essential for implementation. India needs standard documentation formats, institutional pathways, and legal protections for clinicians acting in good faith. These frameworks should be usable not only in large hospitals but also in district and community settings.

Uniformity will reduce confusion and variation between institutions.

Standard forms can help clinicians document prognosis, family discussions, patient preferences, and care plans clearly. Such tools also reassure families that decisions are being made according to accepted norms rather than personal opinion. A national framework can bring confidence to both doctors and the public.

#### **Building an End-of-Life Care Pathway**

Patients with irreversible illness, advanced organ failure, severe neurological injury, terminal malignancy, advanced dementia, or progressive multi-organ dysfunction should be identified early for goals-of-care discussions. Early identification allows time for planning rather than crisis-driven decision-making. Clinicians should recognize warning signs such as repeated ICU admissions, poor functional status, treatment failure, progressive decline, and absence of realistic curative options. Once such patients are identified, communication should begin gradually and sensitively. Waiting until the last hours often deprives families of meaningful participation.

Hospitals should develop structured family meeting processes. These meetings should include explanation of diagnosis, prognosis, treatment options, likely outcomes, and comfort-focused alternatives. A structured meeting helps ensure that important information is not missed. Such meetings should be conducted in a private and calm environment. The family should be allowed to ask questions and express emotions. A single meeting may not be enough; difficult decisions often require repeated conversations and time for acceptance [34].

Difficult cases should have access to ethics consultation and palliative care teams. This provides support to both clinicians and families. Ethics support is especially useful

when there is disagreement, uncertainty, or concern about the appropriateness of continued treatment. Palliative care teams can help manage symptoms and guide communication. Ethics teams can help clarify values, responsibilities, and processes. Together, they reduce conflict and improve confidence in decision-making.

Reassessment also shows families that decisions are not rushed or predetermined [34-39]. It demonstrates that the medical team remains attentive and thoughtful. This ongoing review builds trust and ensures that care remains appropriate to the patient's condition.

#### **Conclusion**

Medicine is not solely about defeating death. It is about honouring life in its entirety, including its final phase. To care well for the dying is not to surrender. It is to uphold the highest ideals of the profession. When cure is no longer possible, dignity must remain non-negotiable.

End-of-life care asks medicine to combine science with wisdom, law with compassion, and technology with humanity. It reminds us that the true measure of healthcare is not only how many lives we save, but also how gently, honestly, and respectfully we care for those who cannot be saved. A dignified death is not a failure of medicine. It is one of its most profound responsibilities. End-of-life care must therefore become a visible, taught, practiced, and protected part of modern healthcare.

The final goal is not only to help people live well, but also to ensure that when life reaches its natural end, patients are not left to suffer unnecessarily. They must be cared for with comfort, honesty, presence, and respect. In that commitment lies the deepest humanity of medicine.

---

**Acknowledgement:** I acknowledge the use of Microsoft Copilot to summarise my initial notes and to proofread my final draft. I also acknowledge the use of other AI tools to make my illustration.

**Conflict of interest:** None to declare.

**Funding:** None

**How to cite:** Javeri Y. End-of-Life Care: Beyond Survival, Towards Dignity. *Ann Med Med Ethics*. 2026;1(1): 5–11. doi.org/10.66765/amme.2026.003

**Submission received:** 8th April 2026

**Submission accepted:** 7th May 2026

**Submission published:** xx.xx.xxxx

#### **ORCID**

Yash Javeri  <http://orcid.org/0000-0002-7384-3637>

**Copyright and license :** All articles in *Annals of Medicine and Medical Ethics* are published under the Creative Commons Attribution–Non Commercial 4.0 International License (CC BY-NC 4.0). This permits unrestricted non-

commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are properly credited.

### References

1. Cassell EJ. The nature of suffering and the goals of medicine. Oxford University Press, USA; 1991 Oct 3.
2. Chochinov HM. Dignity-conserving care—a new model for palliative care: helping the patient feel valued. *Jama*. 2002 May 1;287(17):2253-60.
3. Rome RB, Luminais HH, Bourgeois DA, Blais CM. The role of palliative care at the end of life. *Ochsner journal*. 2011 Dec 21;11(4):348-52.
4. Harstäde CW, Blomberg K, Benzein E, Östlund U. Dignity-conserving care actions in palliative care: An integrative review of Swedish research. *Scandinavian Journal of caring sciences*. 2018 Mar;32(1):8-23.
5. Kennedy G. The importance of patient dignity in care at the end of life. *The Ulster medical journal*. 2016 Jan;85(1):45.
6. Reichlin M. On the ethics of withholding and withdrawing medical treatment. *Multidisciplinary respiratory medicine*. 2014 Jul 16;9(1):39.
7. Kasman DL. When is medical treatment futile? A guide for students, residents, and physicians. *Journal of general internal medicine*. 2004 Oct;19(10):1053-6.
8. Lo JJ, Graves N, Chee JH, Hildon ZJ. A systematic review defining non-beneficial and inappropriate end-of-life treatment in patients with non-cancer diagnoses: theoretical development for multi-stakeholder intervention design in acute care settings. *BMC palliative care*. 2022 Nov 9;21(1):195.
9. Bosslet GT, Pope TM, Rubinfeld GD, Lo B, Truog RD, Rushton CH, Curtis JR, Ford DW, Osborne M, Misak C, Au DH. An official ATS/ AACN/ ACCP/ ESICM/ SCCM policy statement: responding to requests for potentially inappropriate treatments in intensive care units. *American journal of respiratory and critical care medicine*. 2015 Jun 1;191(11):1318-30.
10. Kon AA, Shepard EK, Sederstrom NO, Swoboda SM, Marshall MF, Birriel B, Rincon F. Defining futile and potentially inappropriate interventions: a policy statement from the Society of Critical Care Medicine Ethics Committee. *Critical care medicine*. 2016 Sep 1;44(9):1769-74.
11. Puntillo K, Nelson JE, Weissman D, Curtis R, Weiss S, Frontera J, Gabriel M, Hays R, Lustbader D, Mosenthal A, Mulkerin C. Palliative care in the ICU: relief of pain, dyspnea, and thirst—a report from the IPAL-ICU Advisory Board. *Intensive care medicine*. 2014 Feb;40(2):235-48.
12. Del Río MI, Shand B, Bonati P, Palma A, Maldonado A, Taboada P, Nervi F. Hydration and nutrition at the end of life: a systematic review of emotional impact, perceptions, and decision-making among patients, family, and health care staff. *Psycho-Oncology*. 2012 Sep;21(9):913-21.
13. Carter AN. To what extent does clinically assisted nutrition and hydration have a role in the care of dying people?. *Journal of Palliative Care*. 2020 Oct;35(4):209-16.
14. Tsaloglidou A, Rammos K, Kyparos A, Dimitiadou A, Matziari C. Ethical issues in withholding or withdrawal of artificial nutrition and hydration. *International Journal of Caring Sciences*. 2008 May 1;1(2):66.
15. Brody H, Hermer LD, Scott LD, Grumbles LL, Kutac JE, McCammon SD. Artificial nutrition and hydration: the evolution of ethics, evidence, and policy. *Journal of general internal medicine*. 2011 Sep;26(9):1053-8.
16. Aghabarary M, Nayeri ND. Medical futility and its challenges: a review study. *Journal of medical ethics and history of medicine*. 2016 Oct 20;9:11.
17. Kopar PK, Visani A, Squirrell K, Brown DE. Addressing futility: a practical approach. *Critical care explorations*. 2022 Jul 1;4(7):e0706.
18. da Silva Vieira JV, Deodato S, Mendes F. The concept of futility in health: A scoping review. *Clinical Ethics*. 2021 Dec;16(4):347-53.
19. Bernat JL. Medical futility: definition, determination, and disputes in critical care. *Neurocritical Care*. 2005 Apr;2(2):198-205.
20. Schneiderman LJ, Jecker NS, Jonsen AR. Medical futility: its meaning and ethical implications. *Annals of internal medicine*. 1990 Jun 15;112(12):949-54.
21. Schneiderman LJ. Defining medical futility and improving medical care. *Journal of bioethical inquiry*. 2011 Jun;8(2):123-31.
22. Close E, White BP, Willmott L, Gallois C, Parker M, Graves N, Winch S. Doctors' perceptions of how resource limitations relate to futility in end-of-life decision making: a qualitative analysis. *Journal of Medical Ethics*. 2019 Jun 1;45(6):373-9.
23. Meltzer LS, Huckabay LM. Critical care nurses' perceptions of futile care and its effect on burnout. *American journal of critical care*. 2004 May 1;13(3):202-8.
24. Salins N, Gursahani R, Mathur R, Iyer S, Macaden S, Simha N, Mani RK, Rajagopal MR. Definition of terms used in limitation of treatment and providing

- palliative care at the end of life: The Indian council of medical research commission report. *Indian journal of critical care medicine: peer-reviewed, official publication of Indian Society of Critical Care Medicine*. 2018 Apr;22(4):249.
25. Akdeniz M, Yardımcı B, Kavukcu E. Ethical considerations at the end-of-life care. *SAGE open medicine*. 2021 Mar;9:20503121211000918.
  26. Vincent JL. Withdrawing may be preferable to withholding. *Critical Care*. 2005 Mar 4;9(3):226.
  27. Manalo MF. End-of-life decisions about withholding or withdrawing therapy: medical, ethical, and religious-cultural considerations. *Palliative Care: Research and Treatment*. 2013 Jan;7:PCRT-S10796.
  28. Epner DE, Ravi V, Baile WF. When patients and families feel abandoned. *Supportive Care in Cancer*. 2011 Nov;19(11):1713-7.
  29. Myatra SN, Salins N, Iyer S, Macaden SC, Divatia JV, Muckaden M, Kulkarni P, Simha S, Mani RK. End-of-life care policy: An integrated care plan for the dying: A Joint Position Statement of the Indian Society of Critical Care Medicine (ISCCM) and the Indian Association of Palliative Care (IAPC). *Indian journal of critical care medicine: peer-reviewed, official publication of Indian Society of Critical Care Medicine*. 2014 Sep;18(9):615.
  30. Head BA, Schapmire TJ, Earnshaw L, Chenault J, Pfeifer M, Sawning S, Shaw MA. Improving medical graduates' training in palliative care: advancing education and practice. *Advances in medical education and practice*. 2016 Feb 24:99-113.
  31. Billings JA, Block S. Palliative care in undergraduate medical education: status report and future directions. *Jama*. 1997 Sep 3;278(9):733-8.
  32. VeqAR Z. The perspectives on including palliative care in the Indian undergraduate physiotherapy curriculum. *Journal of Clinical and Diagnostic Research: JCDR*. 2013 Apr 1;7(4):782.
  33. Graham-Wisener L, Nelson A, Byrne A, Islam I, Harrison C, Geddis J, Berry E. Understanding public attitudes to death talk and advance care planning in Northern Ireland using health behaviour change theory: a qualitative study. *BMC Public Health*. 2022 May 6;22(1):906.
  34. Curtis JR, White DB. Practical guidance for evidence-based ICU family conferences. *Chest*. 2008 Oct 1;134(4):835-43.
  35. Kruser JM, Ashana DC, Courtright KR, Kross EK, Neville TH, Rubin E, Schenker Y, Sullivan DR, Thornton JD, Viglianti EM, Costa DK. Defining the time-limited trial for patients with critical illness: an official American Thoracic Society workshop report. *Annals of the American Thoracic Society*. 2024 Feb;21(2):187-99.
  36. Kruser JM, Nadig NR, Viglianti EM, Clapp JT, Secunda KE, Halpern SD. Time-limited trials for patients with critical illness: a review of the literature. *Chest*. 2024 Apr 1;165(4):881-91.
  37. Kesecioglu J, Rusinova K, Alampi D, Arabi YM, Benbenishty J, Benoit D, Boulanger C, Cecconi M, Cox C, Van Dam M, Van Dijk D. European Society of Intensive Care Medicine guidelines on end of life and palliative care in the intensive care unit. *Intensive Care Medicine*. 2024 Nov;50(11):1740-66.
  38. Macaden SC, Salins N, Muckaden M, Kulkarni P, Joad A, Nirabhawane V, Simha S. End of life care policy for the dying: Consensus position statement of Indian association of palliative care. *Indian Journal of Palliative Care*. 2014 Sep;20(3):171.
  39. Gries CJ, Curtis JR, Wall RJ, Engelberg RA. Family member satisfaction with end-of-life decision making in the ICU. *Chest*. 2008 Mar 1;133(3):704-12.